


Croup – Emergency Department

This best practice algorithm provides guidance on diagnosis and medical management for pediatric croup.

Created: July 2026 | Next Review: July 2029

Created by: S. Holmstrom, MD; A. Kenneally, PharmD; Stefan Malin, MD

Reviewed by: Pediatric Critical Care, PEMCET



References:

- Udoh I, Heegeman D, Ravi S. Retrospective Evaluation of Return Rates in Pediatric Patients Treated With Inhaled Racemic Epinephrine for Croup. *WMJ*. 2022 Apr;121(1):26-29. PMID: 35442575.
- Russell KF, Liang Y, O'Gorman K, et al. Glucocorticoids for croup. *Cochrane Database Syst Rev*. 2011;19(1):1-81.
- Jessica Retzke; Croup (Acute Laryngotracheobronchitis). *Quick References* 2021; 10.1542/aap.ppcqr.396247
- Kelly R. Bergmann, Brian Lefchak, Amanda Nickel, Shea Lammers, Dave Watson, Gabrielle Z. Hester; Variation in Organizational Clinical Practice Guidelines for Croup. *Hosp Pediatr* September 2023; 13 (9): e241–e245.



Riley Children's Health
Indiana University Health

CONTACT RILEY CHILDREN'S HEALTH

Calling from an ED or inpatient setting? **Call the IU Health Transfer Center for transfer or consult at 877.247.1177.**

Calling from an outpatient setting? **Call MyRiley at 1.844.697.4539.**

Disclaimer notice: This guideline summarizes publicly available data and is intended for general information purposes only and does not and is not intended to constitute medical treatment. You accept full responsibility for use of information. We do not make any representations to its accuracy, reliability, completeness, timeliness or appropriateness for a particular purpose. We are neither responsible nor liable for any claim, loss or damage resulting from use of this information. The information should not be used in place of a visit, call, consultation or advice of a physician or medical specialist. Information contained herein is not exhaustive and does not cover all diseases, ailments, physical conditions or their treatment. The information contained in this guideline is designed for general use. Each clinician should use their judgment to meet the needs of each individual patient.